

Thomas F. Vail, D.P.M., F.A.C.F.S.

Podiatrist –Foot Surgeon 1725 Western Ave., Suite C Findlay, Ohio 45840 419-423-1888

Board Certified - American Board of Podiatric Surgery

Post-Procedure Instructions for Bleomycin Injection

- Stop for 1 week the versone sal paste and formadon. You may resume these after 2nd week.
- Watch for signs of infections i.e. pain, redness, swelling, pus, red streaking up your foot and or leg, fever or chills. If you should develop any of the above, call our office immediately at 419-423-1888 and tell the person answering the phone that you recently had surgery and the symptoms you are experiencing. The doctor's assistant will let you know the appropriate course of action to take.
- 24-48 hours after procedure remove bandages and bathe as normal. Scrub the surgical site with soap and water. Dry the area and apply Amerigel (preferred and available at our clinic) or Anti-bacterial ointment to the operative site and cover with accommodative dressings.
- Your injection site will be numb for approximately 4 to 5 hours. Begin taking extrastrength Tylenol, Advil, or the prescription written by the Dr. Take the prescription for the first few days whether you have pain or not. Follow the instructions on the label on the bottle. If this does not help with the pain, you may want to call the office at 419-423-1888. If it is after hours the doctor will be paged.

© This form is the property of Dr. Thomas F. Vail, DPM. No portion of this form may be copied or used for commercial, web, print or other unauthorized use. Unauthorized copying of this form for use is strictly forbidden and enforced.